

Discharge Advice Following Angio-Seal

Patient Information

Cardiac Catheter Laboratory
Royal Albert Edward Infirmary

Author ID:	CW
Leaflet Number:	Card 004
Version:	6
Name of Leaflet:	Discharge Advice Following Angio-Seal
Last reviewed:	Jul 2020
Next review date:	Jul 2022

Discharge Advice Following Angio-Seal

- Following your angiogram an Angio-seal device, which is made of collagen has been used to plug the puncture site in your artery.
- The body naturally absorbs the collagen plug in approximately 90 days, during this time carry your patient information card with you at all times.
- You may feel a pea size lump in your groin or mild tenderness at this site.
- You should be accompanied home and have a friend or relative to stay with you over night.
- If severe bleeding occurs at the puncture site, lie down and apply firm pressure. Ask your friend/relative to ring 999 and return immediately to your nearest Accident & Emergency Department.
- If you experience fever, bleeding, persistent tenderness or swelling in the groin, redness and or warm to touch, numbness or pain, discharge or rash you must go immediately to your nearest A&E department.
- Modify activities for 48 hours – no straining, lifting greater than 5kg.
- If coughing, sneezing or straining occurs during the first 12 hours, be advised to place your hand firmly over the puncture site to support it.
- It is advisable to drink plenty of fluids, especially water over the next few days; this will assist with flushing the dye out of your body.
- It is advisable to refrain from the consumption of alcohol for 24 hours.

- A dressing should be left in place for 24 hours. Please refrain from bathing for 3 days following your procedure shower or strip wash only.
- You can remove the dressing the following day, gently clean the site with soap and water. Dry the area gently. Cover the site with a clean dressing until the skin heals approximately 3 days.
- Refrain from driving for at least 48 hours following your angiogram.
- Sometimes bruising of the groin, leg or abdomen may occur and a small lump may be present for several weeks. If you feel pain or the lump enlarges see your GP.
- If you take Metformin **do not** take for 48 hours following your angiogram.
- If you take Warfarin; Apixaban; Rivaroxaban or Edoxaban you may restart on the evening following your angiogram unless the Doctor has instructed otherwise.
- Please contact the Cardiac Catheter Lab on 01942 773392 if you have any questions.

Comments, Compliments or Complaints

The Patient Relations/Patient Advice and Liaison Service (PALS) Department provides confidential on the spot advice, information and support to patients, relatives, friends and carers.

Contact Us

Tel: 01942 822376 (Monday to Friday 9am to 4pm)

The Patient Relations/PALS Manager
Wrightington, Wigan and Leigh NHS Foundation Trust
Royal Albert Edward Infirmary
Wigan Lane
Wigan WN1 2NN

Ask 3 Questions

Become more involved in decisions about your healthcare. You may be asked to make choices about your treatment. To begin with, try to make sure you get the answers to three key questions:

1. What are my options?
2. What are the pros and cons of each option for me?
3. How do I get support to help me make a decision that is right for me?



How We Use Your Information

For details on how we collect, use and store the information we hold about you, please take a look at our “how we use your information” leaflet which can be found on the Trust website: https://www.wwl.nhs.uk/patient_information/leaflets/

This leaflet is also available in audio, large print, Braille and other languages upon request. For more information please ask in the department/ward.

© Wrightington, Wigan and Leigh NHS Foundation Trust
All rights reserved. Not to be reproduced in whole or in part without the permission of the copyright owner

